arXiv:1510.01492v1 [nlin.CD] 6 Oct 2015

Lyapunov dimension formula of attractors in the Tigan and Yang systems

G. A. Leonov, N. V. Kuznetsov, N. A. Korzhemanova, D. V. Kusakin

[a] *Faculty of Mathematics and Mechanics, St. Petersburg State University, 198504 Peterhof, St. Petersburg, Russia*
[b] *Department of Mathematical Information Technology, University of Jyväskylä, 40014 Jyväskylä, Finland*

Abstract

In latter days the technique of attractors dimension estimate of Lorenz type systems is actively developed. In this work the Lyapunov dimension of attractors of the Tigan and Yang systems is estimated.

Keywords: Lorenz-like system, Lorenz system, Yang system, Tigan system, Kaplan-Yorke dimension, Lyapunov dimension, Lyapunov exponents.

1. Introduction

Nowadays various estimates of Lyapunov dimension of attractors of generalized Lorenz systems

$$\begin{cases} \dot{x} = \sigma(y - x), \\ \dot{y} = rx - dy - xz, \\ \dot{z} = -bz + xy, \end{cases} \tag{1}$$

are actively developed [1, 2]. One particular case of (1) system is the Yang system [3]:

$$\begin{cases} \dot{x} = \sigma(y - x), \\ \dot{y} = rx - xz, \\ \dot{z} = -bz + xy, \end{cases} \tag{2}$$

Email address: nkuznetsov239@gmail.com (N. V. Kuznetsov)

where σ, b are positive, r is arbitrary real number. In [4] an estimation of Lyapunov dimension for the generalized Lorenz system (1) was obtained (see also remarks in [5]), which for the case $d = 0$ (i.e. for the Yang system (2)) is as follows:

Theorem 1. *[4] Let K be an invariant compact set of the Yang system (2). Assume $\sigma > 0$, $b > 0$, $r > 0$, $b(\sigma - b) + \sigma r > 0$, and if one of the following conditions holds*

1. $2\sigma \geq b$, $\sigma^2(3b + r) - b^3 - \sigma b(6b + r) > 0$
2. $2\sigma < b$, $3\sigma^2(b + r) - 6\sigma b^2 - b^3 > 0$,

then $\dim_L K \leq 3 - \frac{2(\sigma+b)}{\sigma+\sqrt{\sigma^2+4\sigma r}}$.

In our work we extend the domain of parameters, where the above estimation is valid.

2. Lyapunov functions in the dimension theory

Consider an autonomous differential equation

$$\dot{\mathrm{x}} = \mathrm{f}(\mathrm{x}), \quad \mathrm{x} \in \mathbb{R}^n, \tag{3}$$

where $\mathrm{f} : \mathbb{R}^n \to \mathbb{R}^n$ is smooth. Suppose that solution $\mathrm{x}(t, \mathrm{x}_0)$, $x(0, x_0) = x_0$ exist for all $t \geq 0$ and consider corresponding linearized system along the solution:

$$\dot{\mathrm{u}} = J(\mathrm{x}(t, \mathrm{x}_0))\,\mathrm{u}, \quad \mathrm{u} \in \mathbb{R}^n, \ t \in \mathbb{R}_+, \tag{4}$$

where

$$J(\mathrm{x}(t, \mathrm{x}_0)) = \left[\frac{\partial f_i(\mathrm{x})}{\partial \mathrm{x}_j}\Big|_{\mathrm{x}=\mathrm{x}(t,\mathrm{x}_0)}\right]$$

is the $(n \times n)$ Jacobian matrix. By a linear variable change $\mathrm{y} = S\mathrm{x}$ with a nonsingular $(n \times n)$-matrix S system (3) is transformed into

$$\dot{\mathrm{y}} = S\,\dot{\mathrm{x}} = S\,\mathrm{f}(S^{-1}\mathrm{y}) = \tilde{\mathrm{f}}(\mathrm{y}).$$

Consider the linearization along the corresponding solution $\mathrm{y}(t, \mathrm{y}_0) = S\mathrm{x}(t, S^{-1}\mathrm{x}_0)$:

$$\dot{\mathrm{v}} = \tilde{J}(\mathrm{y}(t, \mathrm{y}_0))\,\mathrm{v}, \quad \mathrm{v} \in \mathbb{R}^n. \tag{5}$$

Here the Jacobian matrix is as follows

$$\tilde{J}(\mathrm{y}(t, \mathrm{y}_0)) = S\,J(\mathrm{x}(t, \mathrm{x}_0))\,S^{-1}. \tag{6}$$

Suppose that $\lambda_1(\mathrm{x}, S) \geqslant \cdots \geqslant \lambda_n(\mathrm{x}, S)$ are eigenvalues of the matrix

$$\frac{1}{2}\left(S\,J(\mathrm{x}(t,\mathrm{x}_0))\,S^{-1} + (S\,J(\mathrm{x}(t,\mathrm{x}_0))\,S^{-1})^*\right). \tag{7}$$

Theorem 2 ([1, 6])**.** *Given an integer $j \in [1, n-1]$ and $s \in [0,1]$, suppose that there are a continuously differentiable scalar function $\vartheta : \mathbb{R}^n \to \mathbb{R}$ and a nonsingular matrix S such that*

$$\lambda_1(\mathrm{x}, S) + \cdots + \lambda_j(\mathrm{x}, S) + s\lambda_{j+1}(\mathrm{x}, S) + \dot{\vartheta}(\mathrm{x}) < 0,\ \forall\, \mathrm{x} \in K. \tag{8}$$

Then $\dim_L K \leqslant j + s$.

Here $\dot{\vartheta}$ is the derivative of ϑ with respect to the vector field f:

$$\dot{\vartheta}(\mathrm{x}) = (\mathrm{grad}(\vartheta))^*\mathrm{f}(\mathrm{x}).$$

The introduction of the matrix S can be regarded as a change of the space metric.

Theorem 3 ([1, 7–9])**.** *Assume that there are a continuously differentiable scalar function ϑ and a nonsingular matrix S such that*

$$\lambda_1(\mathrm{x}, S) + \lambda_2(\mathrm{x}, S) + \dot{\vartheta}(\mathrm{x}) < 0,\ \forall\, \mathrm{x} \in \mathbb{R}^n. \tag{9}$$

Then any solution of system (3) *bounded on* $[0, +\infty)$ *tends to an equilibrium as* $t \to +\infty$.

Thus, if (9) holds, then the global attractor of system (3) coincides with its stationary set.

3. Main result

We can prove the following result for the Yang system.

Theorem 4. 1. *Assume $r = 0$ and the following inequalities $b(\sigma - b) > 0$, $\sigma - \frac{(\sigma+b)^2}{4(\sigma-b)} \geq 0$ are satisfied. Then any bounded on $[0; +\infty)$ solution of system* (2) *tends to a certain equilibrium as $t \to +\infty$.*

2. *Assume $r < 0$ and $r\sigma + b(\sigma - b) > 0$. Then any bounded on $[0; +\infty)$ solution of system* (2) *tends to a certain equilibrium as $t \to +\infty$.*

3. *Assume* $r > 0$ *and there are two distinct real roots* $\gamma^{(II)} > \gamma^{(I)}$ *of equation*

$$4br\sigma^2(\gamma+2\sigma-b)^2+16\sigma b\gamma(r\sigma^2+b(\sigma+b)^2-4\sigma(\sigma r+\sigma b-b^2))=0, \quad (10)$$

such that $\gamma^{(II)} > 0$.
In this case

(a) *if*

$$b(b-\sigma) < r\sigma < b(\sigma+b),$$

then any bounded on $[0;+\infty)$ *solution of system* (2) *tends to a certain equilibrium as* $t \to +\infty$.

(b) *if*

$$r\sigma > b(\sigma+b), \quad (11)$$

then

$$\dim_L K = 3 - \frac{2(\sigma+b)}{\sigma+\sqrt{\sigma^2+4\sigma r}}, \quad (12)$$

where $K \supset (0,0,0)$ *is bounded invariant set of system* (2).

For proving Theorem 4 we use Theorems 2 and 3 with the matrix S of the form

$$S = \begin{pmatrix} -\rho^{-1} & 0 & 0 \\ -\frac{b}{\sigma} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and the function

$$\vartheta(x,y,z) = \frac{(1-s)V(x,y,z)}{[\sigma^2+4\sigma r]^{\frac{1}{2}}},$$

where

$$\rho = \frac{\sigma}{\sqrt{\sigma r + b(\sigma-b)}},$$

$$V(x,y,z) = \gamma_4 x^2 + (\sigma\gamma_2+\gamma_3)y^2 + \gamma_3 z^2 - \frac{1}{4\sigma}\gamma_2 x^4 + \gamma_2 x^2 z + \gamma_1\gamma_2 xy - \frac{\sigma}{b}z.$$

The parameters $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ are variable.

Exact value of the Lyapunov dimension is obtained by the comparison of estimate (9) and the Lyapunov dimension at the zero equilibrium point.

We consider classical chaotic parameters of the Yang system: $\sigma = 10, b = \frac{8}{3}, r = 16$ and $\sigma = 35, b = 3, r = 35$. In the first case inequality (11) is as follows

$$160 > \frac{304}{9}$$

and the greater root of corresponding equation (10) is equal

$$\gamma_1^{(II)} = \frac{6358}{135} + \frac{14}{135}\sqrt{178309}.$$

In the second case inequality (11) is as follows

$$1225 > 114$$

and the greater root of (10) is equal

$$\gamma_1^{(II)} = \frac{193391}{1225} + \frac{2}{1225}\sqrt{7665943314}.$$

Thus, classical chaotic parameters satisfy the requirement of Theorem 4 and the formula (12) is valid.

4. T-system

Consider the Tigan system (T-system) [10, 11]:

$$\begin{cases} \dot{x} = a(y - x), \\ \dot{y} = (c - a)x - axz, \\ \dot{z} = -bz + xy. \end{cases} \tag{13}$$

By the transformation

$$x \to \frac{x}{\sqrt{a}}, y \to \frac{y}{\sqrt{a}}, z \to \frac{z}{a}$$

one has

$$\begin{cases} \dot{x} = a(y - x) \\ \dot{y} = (c - a)x - xz \\ \dot{z} = -bz + xy. \end{cases} \tag{14}$$

So the T-system can be transformed to the Yang system with the following parameters $\sigma = a, r = c - a$. Both systems were independently considered in 2008 year. But a particular case of the T-system was considered in 2004 [12].

The T-system has classical chaotic parameters $(a, b, c) = (2.1, 0.6, 30)$. Conditions of Theorem 4 are valid in the case $\sigma = \frac{21}{10}, b = \frac{6}{10}, r = \frac{279}{10}$ and takes the form:

(1) parameter $r = 27.9 > 0$,

(2) inequality (11) is transformed to

$$\frac{5859}{100} > \frac{81}{50},$$

and, thus, is valid,

(3) the roots of the corresponding equation (10) are $\gamma_1^{(I)} = \frac{9883}{1085} - \frac{1}{1085}\sqrt{82416853}$ and $\gamma_1^{(II)} = \frac{9883}{1085} + \frac{1}{1085}\sqrt{82416853}$, and, thus, the greater root is positive.

Therefore, if $K \supset (0, 0, 0)$ is a bounded invariant set of system (13) with the parameters $\sigma = 2.1, b = 0.6, r = 27.9$, then

$$\dim_L K = 3 - \frac{2(\sigma + b)}{\sigma + \sqrt{\sigma^2 + 4\sigma r}} \approx 2.692345984.$$

5. Conclusions

The Yang system (2) by the transformation

$$x \to \sigma x, y \to \sigma y, z \to \sigma z, t \to \frac{t}{\sigma}, \quad \sigma \neq 0 \tag{15}$$

takes the form

$$\begin{cases} \dot{x} = y - x, \\ \dot{y} = \dfrac{r}{\sigma}x - xz, \\ \dot{z} = -\dfrac{b}{\sigma}z + xy. \end{cases} \tag{16}$$

For $\sigma = 0$ the Yang system becomes linear and its dynamics has minor interest. Thus, without loss of generality, one can assume that $\sigma = 1$. Below we compare the domains of parameters in Theorem 4 and Theorem 1 for $\sigma = 1$.

The points $P1$ and $P2$ are the parameters which correspond the classical chaotic self-excited attractors[1] in the Yang system, $P3$ — classical chaotic self-excited attractor in the T-system, $P4, P5$ — parameters, for which we have also found chaotic self-excited attractors by numerical analysis.

[1]An attractor is called a self-excited attractor if its basin of attraction intersects with any open neighborhood of an equilibrium, otherwise it is called a hidden attractor [13–16].



(a) (b)

Figure 1: Domains of parameters for $\sigma = 1$, $r < 15$. (a) Theorem 4. (b) Theorem 1.



(a) (b)

Figure 2: Self-excited attractors in the Yang system: (a) P4 — $\sigma = 1$, b = 0.8, r = 9.1. (b) P5 — $\sigma = 1$, b = 0.8, r = 9.7.

Up to now in such Lorenz-like systems as Lorenz, Chen, Lu and Tigan systems self-excited chaotic attractors have been found only. In such Lorenz-like systems as Glukhovsky–Dolghansky and Rabinovich systems both self-excited and hidden attractors can be found [16–18]. Recent examples of hidden attractors can be found in *The European Physical Journal Special Topics: Multistability: Uncovering Hidden Attractors*, 2015 (see [19–30]). Note that while coexisting self-excited attractors can be found by the standard computational procedure, there is no regular way to predict the existence or coexistence of hidden attractors.

References

[1] G. A. Leonov, Lyapunov functions in the attractors dimension theory, Journal of Applied Mathematics and Mechanics 76 (2) (2012) 129–141.

[2] G. A. Leonov, N. V. Kuznetsov, On differences and similarities in the analysis of Lorenz, Chen, and Lu systems, Applied Mathematics and Computation 256 (2015) 334–343. doi:10.1016/j.amc.2014.12.132.

[3] Q. Yang, G. Chen, A chaotic system with one saddle and two stable node-foci, International Journal of Bifurcation and Chaos 18 (2008) 1393–1414. doi:10.1142/S0218127408021063.

[4] Y. Chen, Q. Yang, The nonequivalence and dimension formula for attractors of Lorenz-type systems, International Journal of Bifurcation and Chaos 23 (12), art. num. 1350200. doi:10.1142/S0218127413502003.

[5] G. Leonov, N. Kuznetsov, N. Korzhemanova, D. Kusakin, Estimation of Lyapunov dimension for the Chen and Lu systems, arXiv http://arxiv.org/pdf/1504.04726v1.pdf.

[6] G. Leonov, Lyapunov dimension formulas for Henon and Lorenz attractors, St.Petersburg Mathematical Journal 13 (3) (2002) 453–464.

[7] G. A. Leonov, On estimations of the Hausdorff dimension of attractors, Vestnik St.Petersburg University, Mathematics 24 (3) (1991) 41–44.

[8] G. A. Leonov, V. A. Boichenko, Lyapunov's direct method in the estimation of the Hausdorff dimension of attractors, Acta Applicandae Mathematicae 26 (1) (1992) 1–60.

[9] V. A. Boichenko, G. A. Leonov, V. Reitmann, Dimension theory for ordinary differential equations, Teubner, Stuttgart, 2005.

[10] G. Tigan, D. Opris, Analysis of a 3d chaotic system, Chaos, Solitons & Fractals 36 (5) (2008) 1315–1319.

[11] B. Jiang, X. Han, Q. Bi, Hopf bifurcation analysis in the T system, Nonlinear Analysis: Real World Applications 11 (1) (2010) 522 – 527.

[12] G. Tigan, Bifurcation and stability in a system derived from the lorenz system, in: Proceedings of the 3rd. International Colloquium "Mathematics in Engineering and Numerical Physics", October 7-9, 2004: [mathematics Sections], BSG Proceedings, Geometry Balkan Press, 2004, pp. 265–272.

[13] G. A. Leonov, N. V. Kuznetsov, V. I. Vagaitsev, Localization of hidden Chua's attractors, Physics Letters A 375 (23) (2011) 2230–2233. doi:10.1016/j.physleta.2011.04.037.

[14] G. A. Leonov, N. V. Kuznetsov, V. I. Vagaitsev, Hidden attractor in smooth Chua systems, Physica D: Nonlinear Phenomena 241 (18) (2012) 1482–1486. doi:10.1016/j.physd.2012.05.016.

[15] G. A. Leonov, N. V. Kuznetsov, Hidden attractors in dynamical systems. From hidden oscillations in Hilbert-Kolmogorov, Aizerman, and Kalman problems to hidden chaotic attractors in Chua circuits, International Journal of Bifurcation and Chaos 23 (1), art. no. 1330002. doi:10.1142/S0218127413300024.

[16] G. Leonov, N. Kuznetsov, T. Mokaev, Homoclinic orbits, and self-excited and hidden attractors in a Lorenz-like system describing convective fluid motion, Eur. Phys. J. Special Topics 224 (8) (2015) 1421–1458. doi:10.1140/epjst/e2015-02470-3.

[17] N. Kuznetsov, G. A. Leonov, T. N. Mokaev, Hidden attractor in the Rabinovich system, arXiv:1504.04723v1Http://arxiv.org/pdf/1504.04723v1.pdf.

[18] G. Leonov, N. Kuznetsov, T. Mokaev, Homoclinic orbit and hidden attractor in the Lorenz-like system describing the fluid convection motion in the rotating cavity, Communications in Nonlinear Science and Numerical Simulation 28 (doi:10.1016/j.cnsns.2015.04.007) (2015) 166–174.

[19] M. Shahzad, V.-T. Pham, M. Ahmad, S. Jafari, F. Hadaeghi, Synchronization and circuit design of a chaotic system with coexisting hidden attractors, European Physical Journal: Special Topics 224 (8) (2015) 1637–1652.

[20] S. Brezetskyi, D. Dudkowski, T. Kapitaniak, Rare and hidden attractors in Van der Pol-Duffing oscillators, European Physical Journal: Special Topics 224 (8) (2015) 1459–1467.

[21] S. Jafari, J. Sprott, F. Nazarimehr, Recent new examples of hidden attractors, European Physical Journal: Special Topics 224 (8) (2015) 1469–1476.

[22] Z. Zhusubaliyev, E. Mosekilde, A. Churilov, A. Medvedev, Multistability and hidden attractors in an impulsive Goodwin oscillator with time delay, European Physical Journal: Special Topics 224 (8) (2015) 1519–1539.

[23] P. Saha, D. Saha, A. Ray, A. Chowdhury, Memristive non-linear system and hidden attractor, European Physical Journal: Special Topics 224 (8) (2015) 1563–1574.

[24] V. Semenov, I. Korneev, P. Arinushkin, G. Strelkova, T. Vadivasova, V. Anishchenko, Numerical and experimental studies of attractors in memristor-based Chua's oscillator with a line of equilibria. Noise-induced effects, European Physical Journal: Special Topics 224 (8) (2015) 1553–1561.

[25] Y. Feng, Z. Wei, Delayed feedback control and bifurcation analysis of the generalized Sprott B system with hidden attractors, European Physical Journal: Special Topics 224 (8) (2015) 1619–1636.

[26] C. Li, W. Hu, J. Sprott, X. Wang, Multistability in symmetric chaotic systems, European Physical Journal: Special Topics 224 (8) (2015) 1493–1506.

[27] Y. Feng, J. Pu, Z. Wei, Switched generalized function projective synchronization of two hyperchaotic systems with hidden attractors, European Physical Journal: Special Topics 224 (8) (2015) 1593–1604.

[28] J. Sprott, Strange attractors with various equilibrium types, European Physical Journal: Special Topics 224 (8) (2015) 1409–1419.

[29] V. Pham, S. Vaidyanathan, C. Volos, S. Jafari, Hidden attractors in a chaotic system with an exponential nonlinear term, European Physical Journal: Special Topics 224 (8) (2015) 1507–1517.

[30] S. Vaidyanathan, V.-T. Pham, C. Volos, A 5-D hyperchaotic Rikitake dynamo system with hidden attractors, European Physical Journal: Special Topics 224 (8) (2015) 1575–1592.